



Authorization criteria: Knee arthroplasty for adults, total, revision

**For BCN HMOSM (commercial) and BCN AdvantageSM members
For Blue Cross Medicare Plus BlueSM PPO members**

Clinical review is required for adult members (age ≥18). The medical necessity criteria are outlined below.

CPT codes: *27486, *27487, *27488

Fractured prosthesis or cement

The following must be met:

1. Fractured prosthesis or cement identified by imaging
2. No active infection in any location

Malposition of tibial or femoral component

The following must be met:

1. Malposition of tibial or femoral component identified by imaging
2. No active infection in any location

Recurrent dislocation

The following must be met:

1. Recurrent dislocation identified by imaging
2. No active infection in any location

Symptomatic loosening of prosthesis or cement

The following must be met:

1. Symptomatic loosening of prosthesis or cement identified by imaging
2. No active infection in any location

Worn or dislocated plastic insert

The following must be met:

1. Worn or dislocated plastic insert identified by imaging
2. No active infection in any location



Authorization criteria: Knee arthroplasty for adults, total, revision

**For BCN HMOSM (commercial) and BCN AdvantageSM members
For Blue Cross Medicare Plus BlueSM PPO members**

Joint infection

Both of the following must be met:

1. Joint infection indicated by one or more of the following:
 - a. Sinus tract communicating with prosthetic joint identified by imaging
 - b. Prosthetic joint infection by positive synovial fluid culture or tissue culture with ONE OR MORE of the following:
 - i. Two cultures positive for the same organism
 - ii. Culture positive for Staphylococcus aureus (S. aureus)
 - iii. Culture positive for gram negative organism
 - iv. Culture positive for enterococci
 - c. Signs and symptoms of infection as indicated by BOTH of the following:
 - i. Prosthetic joint signs and symptoms of ONE OR MORE of the following:
 1. Joint pain, OR
 2. Erythema, OR
 3. Drainage at the joint on physical exam, OR
 4. Swelling at the joint by physical exam
 - ii. Findings of TWO OR MORE of the following:
 1. Temperature >100.4 (38.0 C)
 2. Synovial WBC or neutrophil percentage >normal
 3. ESR >30 mm/hr
 4. C-reactive protein >normal
2. BOTH of the following:
 - a. EITHER of the following:
 - i. Joint infection within the immediate post-operative period of 4 weeks from total joint replacement, OR
 - ii. New joint symptoms and findings after the immediate postoperative period that are greater than 3 weeks in duration
 - b. EITHER of the following:
 - i. Loosening of prosthesis or cement identified by imaging, OR
 - ii. Continued symptoms or findings after treatment with BOTH of the following:
 1. IV anti-infectives for at least for 4 weeks, AND
 2. Joint lavage AND drainage

References

McKesson's InterQual[®] 2017 Procedures Adult Criteria Removal and Replacement, Total Joint Replacement (TJR) Knee

*CPT codes, descriptions and two-digit numeric modifiers only are copyright 2017 American Medical Association. All rights reserved.

Current authorization criteria effective date: March 2018